arXiv:cond-mat/0008166v1 [cond-mat.str-el] 10 Aug 2000

Mutagenesis and Metallic DNA

Chun-Min Chang[1], A. H. Castro Neto[1], and A. R. Bishop[2]
[1] *Department of Physics, University of California, Riverside, CA 92521*
[2] *Theoretical Division and Center for Nonlinear Studies, Los Alamos National Laboratory, Los Alamos, New Mexico 87545*

(February 1, 2008)

Positions of protons in DNA hydrogen bonds are fundamental for the fidelity of the replication process. Because the hydrogen bond is partially covalent the electronic structure of DNA plays an important role in the replication process. Current studies of electron transfer in DNA have shown that simple pictures based on independent electron physics are not sufficient to account for the experimental data. In this paper we propose a model for electron-proton interaction in DNA and its relationship to mutagenesis. We argue that mutations will be more frequent in DNA sequences where electron delocalization occurs. To test our theory we calculate the heat capacity of DNA at low temperatures and compare with the available experimental data in hydrated DNA. We find that our model is in good agreement with the data. We also propose new measurements of infrared absorption which would further test our ideas.

PACS numbers: 87.16.U, 87.14.G, 87.15.M

Watson and Crick [1] pointed out the importance of proton transfer as a possible cause of tautomeric base pairs in DNA and its deep relationship to mutagenesis. Errors in the replication process of DNA can lead to the occurrence of mutations related to genetic diseases and some forms of cancer. Löwdin [2] argued that proton transfer is a quantum mechanical effect which has purely electrostatic origin. Since then many researchers have calculated the potential energy profiles for proton transfer reactions in isolated base pairs [3]. These studies have shown the importance of the electronic configuration at each base pair as a stabilizing factor of the hydrogen bond (H-bond). Indeed, recent experiments [4] show the occurrence of proton transfer in excited base pairs. The problem of electron transfer in DNA has been a source of intense debate [5]. Many different experimental studies of electron transfer by chemical reaction have shown that long range electron transfer in DNA may be rather different from the same process in proteins, where single electron processes are dominant [6]. In this paper we propose a model which combines two essential ingredients in the physics of DNA: the electronic degrees of freedom, and the protons in the H-bonds along the double helix. Our model is different from other proposed models for electron-proton interactions in DNA [7], since we take into account (*i*) the quantum mechanical character of electron motion, (*ii*) the charging energy due to electron transfer between different base pairs, and (*iii*) the specificity of the electron-proton processes due to differences in base pair sequences.

Let us consider a periodic DNA sequence. It is well-known that the electrons in such DNA can move by the overlap of π orbitals along the DNA helix [8]. In this case DNA can be viewed as an one-dimensional (1D) solid. Because a periodic DNA provides a periodic potential for electron motion, basic quantum mechanics tells us that the quantum states of the system are Bloch waves with wavenumber k and band index n. To each value of k we associate a wavelength λ of the electron ($k = 2\pi/\lambda$), and for each band we associate a molecular or atomic state. Band structure calculations [9] indicate that such a DNA sequence should be an insulator with a band gap of order of a few electron volts. In this case the low energy excitations of the DNA are excitons. Recent experiments in Poly(G)-Poly(C) DNA seem to confirm this picture [10]. However, electronic states in biologically relevant DNA (non-periodic) depend considerably on the environmental conditions (such as pH, hydration, etc). Although the study of periodic DNA sequences is very important, it probably does not provide final information on the behavior of electrons in DNA under conditions where mutagenesis is relevant.

Recent experiments on long range electron transfer in DNA indicate that the introduction of Guanine-Citosine (G-C) base pairs enhance electron transfer [11]. This situation is analogous to that in semiconductors where the introduction of electron donors (acceptors) increases the charge conduction by the creation of localized states close to the band edges. When a sufficiently large number of donors or acceptors is introduced into the DNA molecule an impurity band can be formed and conduction is possible. This picture seems to be supported by recent measurements of DNA resistivity [12]. Another possible source of 1D conduction is via non-linear excitations such as solitons [13]. As is well-known, disorder in 1D systems leads to localization of electron wavefunctions [14]. However, when the localization length is large compared with typical electronic scales, electron transfer occurs. Moreover, DNA is aperiodic rather than random, and Anderson localization is a weaker effect [15].

We assume from the outset that the charge carriers (electrons or solitons) of DNA are delocalized over a region of size L such that the energy level spacing $\hbar v_{F,l}/L$ (where $v_{F,l}$ is the Fermi velocity in the strand l) is much smaller than the thermal energy $k_B T$ (where k_B is the Boltzmann constant and T is the temperature). The low energy excitations in this case are particle-hole pairs close to the Fermi points at $\pm k_{F,l}$. This simple scenario, however, is not enough to describe charge propagation in DNA because the charging energies for electron transfer between different base pairs is of order of electron volts as it is common in organic systems. From this point of view DNA is very close to being a Mott insulator instead of an ordinary band insulator [14]. In 1D, such a conductor is called a Luttinger liquid [16]. In a Luttinger liquid the low lying excitations are bosonic collective modes of spin and charge which have solitonic character.

Long ago Pauling [17] proposed that H-bonds have important covalent character, which has subsequently been confirmed [18]. Proton states at the H-bonds are very important because they stabilize the double helix. Protons are much heavier than the electrons and therefore much slower. In A-T base pairs there are 2 H-bonds anchoring the two strands of DNA, while in the G-C case there are 3 H-bonds. In principle each proton in the H-bond can occupy one of its sides, and therefore there are 4 (8) possible states for the protons in A-T (C-G) base pairs. These states do not have the same energy because of the electrostatic interaction between the atoms in the base pairs with the protons at the H-bonds. Moreover, the difference in energy between the states is usually much larger than the thermal energy and therefore we only need to consider the ground state and the first excited state. We call the excitation energy between these two states ϵ_m, where m labels the position of the base pair along the DNA backbone. In this case the proton Hamiltonian can be simply written as

$$H_p = -\frac{1}{2}\sum_m \epsilon_m \sigma^z_m \tag{1}$$

where σ^z is a Pauli matrix. In this representation the ground state of the proton problem is the pseudo-spin state $|\Uparrow\rangle$ while the excited state is $|\Downarrow\rangle$. Notice that ϵ_m can have only two values corresponding to a A-T or G-C base pair. Thus, strictly speaking ϵ_m is the energy required to produce a tautomeric state on an isolated base pair. Because of the large proton mass we disregard proton tunneling and treat the proton as a classical particle. Notice that at temperature T the probability of having a local proton configuration in a mutated state is given by the Boltzmann distribution:

$$P_m(T) \propto \exp\left[-\epsilon_m/(k_B T)\right] \tag{2}$$

corresponding to a thermal jump over the energy barrier. At room temperature and with ϵ_m of the order of tenths of eV this probability is very small and mutation is a rare event.

Let us now consider the coupling between protons and π electrons. This coupling depends strongly on the degrees of freedom of the H-bonds and the π orbitals. On the one hand, the protons in the H-bond have their own magnetic spin degrees of freedom and non-magnetic pseudo-spin variables associated with their position in the H-bond. On the other hand, electron states in DNA can be labeled by their momentum $\hbar k$, their true spin magnetic quantum number, and by their two possible strand numbers: $l = \pm 1$. While the strand and the spin quantum numbers of the electrons couple directly to the pseudo-spin of the proton, the proton spin couples to the electron spin via dipolar forces or contact interactions. If we view the single H-bond problem as an impurity in a chain (or ladder) then the DNA problem is of the so-called multi-channel Kondo type [19]. Since in this paper we are only interested with the stability of the H-bond due to the presence of electrons and not with the magnetic properties of DNA, we neglect completely the dipolar interactions which can be studied separately. Moreover, the symmetry of the coupling between pseudo-spins and electrons is reduced due to the strong anisotropy of the proton positions. Therefore, in the simplest model, the electron-proton problem reduces to a local coupling of value $\lambda_{m,l}$ between the electronic charge density and the H-bond at the base pair m on the strand l:

$$H_{e-p} = \sum_{m,l} \lambda_{m,l}\left[\rho_{m,l} - \overline{\rho}_{m,l}\right](\sigma^z_m - 1) \tag{3}$$

where $\overline{\rho}_{m,l}$ is the average electron density of the mth base pair in the lth strand and $\rho_{m,l}$ the actual electronic density. The physics of this problem is rather simple. In the absence of electronic coupling the H-bond of a given base pair m will find itself in the lowest energy state which is stable at room temperature. Because the proton is coupled to the electrons it can generate electronic displacements along the helix with a gain of electrostatic energy. This gain of electrostatic energy, in turn, can stabilize a tautomeric configuration of the protons leading to mutation. The main question is whether the gain in energy due to the electron-proton interaction is enough to compensate for the loss of electrostatic energy between the proton and the base pair. Moreover, the coupling between H-bonds and electrons also generates an exchange interaction between H-bonds, which oscillates in space with a wavelength of order of the distance a between base pairs.

The low energy excitations of this Luttinger liquid can be studied by the bosonization technique in which solitonic modes are bosons with momentum $\hbar k$, energy $\hbar v_{F,l}|k|$ and strand number $l = \pm 1$. In Hamiltonian form it can be written as [16]

$$H_L = \sum_{k,l} \hbar \overline{v}_{F,l}|k| b^\dagger_{k,l} b_{k,l}$$

where $b_{k,l}$ ($b^\dagger_{k,l}$) is the boson annihilation (creation) operator and $\overline{v}_{F,l} = v_{F,l}\sqrt{1 + 2U_l/(\pi v_{F,l})}$ is the renormalized Fermi velocity due to the electron-electron interaction strength U_l. In our case the spin degrees of freedom are completely decoupled (unless we introduce a direct coupling between the proton and the electron spin) and do not appear explicitly into the problem. In particular, the density operator which appears in (3) can be written as

$$\rho_{m,l} = \sum_k k\sqrt{\frac{L}{2\pi |k|}} e^{-\theta_l} \left(b_{k,l} e^{ikx_m} + b^\dagger_{-k,l} e^{-ikx_m} \right)$$

where x_m is the position of the mth pair along the helix and $e^{\theta_l} = [1 + U_l/(\pi v_{F,l})]^{1/4}$. The main gain in using the bosonization procedure for this problem is that it maps the DNA problem given by the Hamiltonian $H = H_p + H_L + H_{e-p}$ into an exactly solvable problem in many-body physics called *the independent boson model* [20]. Our model describes the relaxation of the Luttinger liquid along the helix due to the proton configuration.

The first result of the model is that the excitation energy ϵ_m of the H-bond is reduced by an amount Δ_m, where

$$\Delta_m = \frac{1}{2} \sum_{l=\pm 1} \frac{\lambda^2_{m,l}}{\pi v_{F,l} + U_l} \, . \tag{4}$$

Thus, in the presence of a *local* metallic electron fluid one gets a *reduction* of the “natural” bias in the system and an increase in the probability (2) of finding the system in a tautomeric state. It is clear from (4) that this can only happen when coupling between H-bonds and electrons is large or when the Coulomb interaction between electrons is weak. In metallic DNA the electron-electron interactions will be screened by the electron motion and therefore Δ_m can be large enough to induce mutation at room temperature. This mechanism for mutation seems to be consistent with recent findings relating the existence of repetitive sequences in DNA (which tend to delocalize the electrons) and mutagenesis associated with evolution [21].

In order to test our ideas we have calculated the heat capacity of DNA using the above described model. There are three main contributors to the DNA specific heat at low temperatures: the phonons which give a contribution growing like T^3, the electrons which contribute with a term proportional to T, and the protons which contribute with a Schottky term. The total specific heat can be calculated once the DNA sequence is known. We assume that the renormalized bias $\overline{\epsilon}_m = \epsilon_m - \Delta_m$ can be written as a series expansion of the temperature: $\overline{\epsilon}_m \approx A_m + B_m T$. The temperature dependence of the bias is due to the exchange interaction between the H-bonds propagated by the electron fluid. In a sample of DNA the values of A_m and B_m are distributed around an average value which is given by the average number of G-C and A-T pairs. For simplicity we assume here a gaussian distribution for both quantities. In Fig. 1 we compare the results of our model with the experiments reported in Ref. [22]. The agreement between our model and the experiments is good. However, there are serious questions about sample quality in experiments dealing with DNA and therefore we have also calculated, as a further check, the infrared (IR) absorption for DNA using many-body techniques. The correlation function of interest is the proton spectral function which is defined as $S_m(\omega) = -2Im[G_m(\omega)]$ where $G_m(\omega)$ is the Green's function of the proton at base pair m as a function of the frequency ω. We found that IR absorption at a certain site m is controlled by a parameter g_m defined as

$$g_m = \frac{a\Delta_m}{\pi \hbar \overline{v}_F} \, . \tag{5}$$

At zero temperature and $g_m < 1$, DNA shows an X-ray edge effect with a sharp absorption edge. This effect is a result of the many-body character of DNA electronic conduction. If $g_m > 1$ or the temperature is finite the IR absorption becomes smooth. For a specific DNA sequence one has to calculate the absorption of the DNA as a whole. In Fig. 2 we show the absorption spectrum for one of the DNA sequences of Fig. 1 at finite temperatures. Observe that no sharp X-ray edge is visible because of the superposition of many X-ray edges coming from individual protons. There is, however, a suppression of the absorption at low frequencies with the formation of a “pseudo-gap”.

In conclusion, we have proposed a model for electron-proton coupling and mutagenesis in DNA based on the idea of charge delocalization. We have shown that the position of the protons in DNA can change, leading to mutations when the π electrons delocalize. We have compared our results for the specific heat at low temperatures with the available experimental data and found near quantitative agreement. We have also proposed new IR absorption experiments which can further test our model.

We acknowledge partial support from a Los Alamos CULAR grant under the auspices of the U. S. Department of Energy.

---

[1] Watson, J. D. H. & Crick, F. H. C., Nature **171**, 737-738 (1953).

[2] Löwdin, P.-O., Adv. Quantum Chem. **2**, 213-361 (1965).

[3] Douhal, A. *et al.*, J. M., Chem. Phys. Lett. **256**, 370- 376 (1996).

[4] Douhal, A., Kim, S. K. & Zewail, A. H., Nature **378**, 260-263 (1995); Folmer, D. E. *et al.*, A. W., Chem. Phys. Lett. **287**, 1-7 (1998).

[5] Wilson, E. K. , C&EN **23**, 43 (1999).

[6] Núñez, M. E., Hall, D. B. & Barton, J. K., Chemistry and Biology **6**, 85 (1999).

[7] Steenken, S., Biol. Chem. **378**, 1293-1297 (1997).

[8] Eley, D. D. & Spivey, D. I., Trans. Faraday Soc. **58**, 411 (1962).
[9] Lewis, J. P. *et al.*, Phys. Rev. B **55**, 6880 (1997).
[10] D. Porath *et al.*, Nature **403**, 635 (2000).
[11] Lewis, F. D. *et al.*, Science **277**, 673 (1997), Meggers, E. *et al.*, J. Am. Chem. Soc. **120**, 12950 (1998).
[12] Fink, H.-W. & Schönengerg, C., Nature **398**, 407 (1999).
[13] A. H. Castro Neto and A. O. Caldeira, Phys. Rev. E **48**, 4037 (1993).
[14] Mott, N. F. in *Metal-Insulator transition* (Taylor & Francis, London 1974).
[15] Vida, J. *et al.*, Phys. Rev. Lett. **83**, 3908 (1999).
[16] Voit, J., Rep. Prog. Phys. **58**, 977 (1995).
[17] Pauling, L. in *The nature of the chemical bond* (Cornell Press, Ithaca 1960).
[18] Isaacs, E. D. *et al.*, Phys. Rev. Lett. **82**, 660 (1999).
[19] Nozières, P. & Blandin, A., J. Physique **41**, 193 (1980).
[20] G. Mahan in *Many-Particle Physics* (Plenum Press, New York 1990) pg. 285.
[21] Moxon, E. R. & Wills, C., Scientific American **280**, 94 (1999).
[22] Mrevlishvili, G. M. *et al.*, Biophysics **40**, 518 (1995).

FIG. 1. Specific heat of DNA : 1. DNA sample with water content $n_0 = 0 - 2$ M H_2O/MBP. The results are presented by the continuous curve (theoretical) and $\circ$ (experimental). 2. DNA sample with water content $n_s = 10 - 12$ M H_2O/MBP (mostly A-form DNA). The results are presented by the dotted curve (theoretical) and $\square$ (experimental). 3. DNA sample with water content $n_\Sigma = 22 - 23$ M H_2O/MBP (mostly B-form DNA). The results are presented by the dashed curve (theoretical) and $\triangle$ (experimental).

FIG. 2. Predicted infrared absorption at finite temperatures, $h\bar{v}_F/a = 2\,\mathrm{eV}$.


Amplitude (arbitary units)
4
3.5
3
2.5
2
1.5
1
0.5
0
0.05
0.1
0.15
0.2
0.25
w (eV)
1
2
3
4
5
1 : T = 5 K
2 : T = 25 K
3 : T = 50 K
4 : T = 100 K
5 : T = 150 K


$C_p \times 10^3$ J / (g×K)
8
6
4
2
1
2
3
2
3
4
5
6
7
8
9
T (K)